

MEMO 2023-12ED-00010(2) Geometric series

- (1) Calculate the first three terms of the GS with $S_{\infty} = 54$ and $T_2 = 12$.

$$S_{\infty} = \frac{a}{1-r}$$

$$T_n = a \cdot r^{n-1}$$

$\therefore$

$$54 = \frac{a}{1-r}$$

$$\therefore T_2 = a \cdot r^{2-1}$$

$$\therefore 54(1-r) = a$$

$$\therefore 12 = a \cdot r^1$$

$$\therefore 54 - 54r = \frac{12}{r}$$

$$\therefore \frac{12}{r} = a$$

$$\therefore 54r - 54r^2 = 12$$

$$\therefore 0 = 54r^2 - 54r + 12$$

$$\therefore 0 = 9r^2 - 9r + 2 \quad [\div 6]$$

$$\therefore 0 = (3r-2)(3r-1)$$

$$\therefore r = \frac{2}{3} \quad \text{or}$$

$$r = \frac{1}{3}$$

$$\text{but } a = \frac{12}{r}$$

$$a = \frac{12}{r}$$

$$\therefore a = 12 \div \frac{2}{3}$$

$$a = 12 \div \frac{1}{3}$$

$$\therefore a = 12 \times \frac{3}{2}$$

$$a = 12 \times \frac{3}{1}$$

$$\therefore a = 18$$

$$a = 36$$

$$\therefore \text{Sequence: } 18 ; 12 ; 8$$

$$\therefore \text{Sequence: } 36 ; 12 ; 4$$

- (2) Consider the GS: $(k+2) + (k+2)^2 + (k+2)^3 + \dots$

- (a) For which value(s) of k will the series converge?

Convergent if: $-1 < r < 1 ; r \neq 0$

$$r = (k+2)$$

$$\therefore -1 < k+2 < 1 ; k+2 \neq 0$$

$$\therefore -1-2 < k < 1-2 ; k \neq -2$$

$$\therefore -3 < k < -1 ; k \neq -2$$

- (b) Calculate, in terms of k , the sum to infinity of the series if it is convergent.

$$a = k+2$$

$$S_{\infty} = \frac{a}{1-r}$$

$$r = (k+2)$$

$$\therefore S_{\infty} = \frac{k+2}{1-(k+2)}$$

$$\therefore S_{\infty} = \frac{k+2}{1-k-2}$$

$$\therefore S_{\infty} = \frac{k+2}{-k-1}$$

(3) Calculate:

(a) $2 - 4 + 8 - 16 + \dots + 512$ **GS**

$$a = 2 \qquad T_n = a \cdot r^{n-1}$$

$$r = -2 \qquad \therefore 512 = 2 \times (-2)^{n-1}$$

$$n = ? \qquad \therefore \frac{512}{2} = (-2)^{n-1}$$

$$T_n = 512 \qquad \therefore 256 = (-2)^{n-1}$$

$$\therefore (-2)^8 = (-2)^{n-1}$$

$$\therefore n - 1 = 8 \qquad \rightarrow \qquad \text{EB} \Leftrightarrow \text{EE}$$

$$\therefore n = 9$$

$$a = 2 \qquad S_n = \frac{a(r^n - 1)}{r - 1}$$

$$r = -2 \qquad \therefore S_9 = \frac{2((-2)^9 - 1)}{-2 - 1}$$

$$n = 9 \qquad \therefore S_9 = \frac{2((-2)^9 - 1)}{-3}$$

$$S_n = ? \qquad \therefore S_9 \approx 342$$

(b) $-38 - 35,5 - 33 - 30,5 - \dots - 10,5$ **AS**

$$a = -38 \qquad T_n = a + (n - 1)d$$

$$d = 2,5 \qquad \therefore -10,5 = -38 + (n - 1)(2,5)$$

$$n = ? \qquad \therefore -10,5 + 38 = 2,5n - 2,5$$

$$T_n = -10,5 \qquad \therefore 27,5 + 2,5 = 2,5n$$

$$\therefore 30 = 2,5n$$

$$\therefore n = \frac{30}{2,5}$$

$$\therefore n = 12$$

$$a = -38 \qquad S_n = \frac{n}{2}[2a + (n - 1)d]$$

$$d = 2,5 \qquad \therefore S_{12} = \frac{12}{2}[2 \times (-38) + (12 - 1)(2,5)]$$

$$n = 12 \qquad \therefore S_{12} = 6[-76 + 11 \times 2,5]$$

$$S_n = ? \qquad \therefore S_{12} = -291$$

(c) $\frac{2}{3} + \frac{2}{9} + \frac{2}{27} + \dots$ to seven term **GS**

$$a = \frac{2}{3} \quad S_n = \frac{a(r^n - 1)}{r - 1}$$

$$r = \frac{1}{3} \quad \therefore S_7 = \frac{\frac{2}{3}\left(\left(\frac{1}{3}\right)^7 - 1\right)}{\frac{1}{3} - 1}$$

$$n = 7 \quad \therefore S_7 = \frac{\frac{2}{3}\left(\left(\frac{1}{3}\right)^7 - 1\right)}{-\frac{2}{3}}$$

$$S_n = ? \quad \therefore S_7 = \frac{2186}{2187}$$

(d) $\sum_{n=5}^{12} \left(\frac{1}{2} \times 3^n\right)$

Sequence: $\frac{1}{2} \times 3^5 + \frac{1}{2} \times 3^6 + \frac{1}{2} \times 3^7 + \dots + \frac{1}{2} \times 3^{12} = \frac{243}{2} + \frac{729}{2} + \frac{2187}{2} + \dots$ **8 terms**

$$a = \frac{243}{2} \quad S_n = \frac{a(r^n - 1)}{r - 1}$$

$$r = 3 \quad \therefore S_8 = \frac{\frac{243}{2}((3)^8 - 1)}{3 - 1}$$

$$n = 8 \quad \therefore S_8 = \frac{\frac{243}{2}((3)^8 - 1)}{2}$$

$$S_n = ? \quad \therefore S_8 = 398520$$

(4) The third term of a GS is 3 and the seventh term is $15\frac{3}{16}$. Calculate:

(a) the constant ratio of the sequence.

$$T_3 = 3 \quad T_7 = 15\frac{3}{16}$$

$$\therefore T_3 = a \cdot r^{3-1} \quad \therefore T_7 = a \cdot r^{7-1}$$

$$\therefore 3 = a \cdot r^2 \dots \textcircled{1} \quad \therefore 15\frac{3}{16} = a \cdot r^6 \dots \textcircled{2}$$

$$\therefore \frac{\frac{243}{16}}{3} = \frac{a \cdot r^6}{a \cdot r^2} \quad \textcircled{2} \div \textcircled{1}$$

$$\therefore \frac{81}{16} = \frac{a \cdot r^6}{a \cdot r^2}$$

$$\therefore \frac{81}{16} = r^{6-2}$$

$$\therefore \left(\frac{3}{2}\right)^4 = r^4 \quad \text{but } EB \Leftrightarrow EE$$

$$\therefore r = \pm \frac{3}{2} \rightarrow \pm \text{ because exponent is even}$$

(b) the first term of the sequence.

Substitute r in (1) or (2):

$$\therefore a \cdot r^2 = 3 \quad \dots \dots \textcircled{1}$$

$$\therefore a \cdot \left(\pm \frac{3}{2}\right)^2 = 3$$

$$\therefore a \cdot \frac{9}{4} = 3$$

$$\therefore a = 3 \times \frac{4}{9}$$

$$\therefore a = \frac{4}{3}$$

$$\text{or} \quad 15 \frac{3}{16} = a \cdot r^6 \quad \dots \textcircled{2}$$

$$a \cdot \left(\pm \frac{3}{2}\right)^6 = \frac{243}{16}$$

$$a \cdot \frac{729}{64} = \frac{243}{16}$$

$$\therefore a = \frac{243}{16} \times \frac{64}{729}$$

$$\therefore a = \frac{4}{3}$$

(c) which term is equal to $76 \frac{227}{256}$.

$$a = \frac{4}{3}$$

$$T_n = a \cdot r^{n-1}$$

$$r = \pm \frac{3}{2} \quad \therefore 76 \frac{227}{256} = \frac{4}{3} \times \left(\pm \frac{3}{2}\right)^{n-1}$$

$$n = ? \quad \therefore \frac{19\,683}{256} = \frac{4}{3} \times \left(\pm \frac{3}{2}\right)^{n-1}$$

$$T_n = 76 \frac{227}{256} \quad \therefore \frac{19\,683}{256} \times \frac{3}{4} = \left(\pm \frac{3}{2}\right)^{n-1}$$

$$\therefore \frac{59\,049}{1\,024} = \left(\pm \frac{3}{2}\right)^{n-1}$$

$$\therefore \left(\pm \frac{3}{2}\right)^{10} = \left(\pm \frac{3}{2}\right)^{n-1} \quad \text{but } EB \Leftrightarrow EE$$

$$\therefore 10 = n - 1$$

$$\therefore n = 11$$

(5) Calculate n if $3 + 12 + 48 + 192 + \dots +$ to n terms $= 1\,048\,575$

$$a = 3 \quad S_n = \frac{a(r^n - 1)}{r - 1}$$

$$r = 4 \quad \therefore 1\,048\,575 = \frac{3(4^n - 1)}{4 - 1}$$

$$n = ? \quad \therefore 1\,048\,575 = \frac{3(4^n - 1)}{3}$$

$$S_n = 1\,048\,575 \quad \therefore 1\,048\,575 = \frac{3(4^n - 1)}{3}$$

$$\therefore 1\,048\,575 = 4^n - 1$$

$$\therefore 1\,048\,576 = 4^n$$

$$\therefore 4^{10} = 4^n \quad \text{but } EB \Leftrightarrow EE$$

$$\therefore n = 10$$

(6) Consider the series: $384 + 192 + 96 + 48 + \dots$

(a) Write the series in sigma notation.

$$\begin{aligned} a &= 384 & T_n &= a \cdot r^{n-1} \\ r &= \frac{1}{2} & \therefore T_n &= 384 \cdot \left(\frac{1}{2}\right)^{n-1} \\ n &=? & \therefore T_n &= 384 \cdot (2^{-1})^{n-1} \\ T_n &=? & \therefore T_n &= 384 \cdot (2)^{1-n} \end{aligned}$$

$$\therefore \sum_{n=1}^k (384 \times 2^{1-n})$$

(b) Determine S_∞ , if possible.

$$\begin{aligned} S_\infty &= \frac{a}{1-r} \\ \therefore S_\infty &= \frac{384}{1-\frac{1}{2}} \end{aligned}$$

$$\therefore S_\infty = 768$$

(c) Calculate T_{12} .

$$\begin{aligned} T_n &= 384 \cdot (2)^{1-n} & \rightarrow & \text{ sien (a)} \\ \therefore T_{12} &= 384 \cdot (2)^{1-12} \\ &= 384 \cdot (2)^{-11} \\ \therefore T_{12} &= \frac{3}{16} \end{aligned}$$

(d) Determine how many terms will have a sum of 762.

$$\begin{aligned} a &= 384 & S_n &= \frac{a(r^n - 1)}{r - 1} \\ r &= \frac{1}{2} & \therefore 762 &= \frac{384\left(\left(\frac{1}{2}\right)^n - 1\right)}{\frac{1}{2} - 1} \\ n &=? & \therefore 762 &= \frac{384\left(\left(\frac{1}{2}\right)^n - 1\right)}{-\frac{1}{2}} \\ S_n &= 762 & \therefore 762 &= -768\left(\left(\frac{1}{2}\right)^n - 1\right) \\ & & \therefore \frac{762}{-768} &= \left(\frac{1}{2}\right)^n - 1 \\ & & \therefore -\frac{127}{128} &= \left(\frac{1}{2}\right)^n - 1 \\ & & \therefore 1 - \frac{127}{128} &= \left(\frac{1}{2}\right)^n \\ & & \therefore \frac{1}{128} &= \left(\frac{1}{2}\right)^n \\ & & \therefore \left(\frac{1}{2}\right)^7 &= \left(\frac{1}{2}\right)^n & \text{ but } EB \Leftrightarrow EE \\ & & \therefore n &= 7 \end{aligned}$$