


The graphs of $f(x) = 2^x - 8$ and $g(x) = ax^2 + bx + c$ sketched below. B and C(0 ; 4,5) are the y-intercepts of the graphs of f and g respectively. The two graphs intersect at A, which is the turning point of the graph of g and the x-intercept of the graphs of f and g .



- (1) Determine the coordinates of A and B.

(4)

A: x-intercept of $g \rightarrow y = 0$

B: y-intercept of $g \rightarrow x = 0$

$$\rightarrow 0 = 2^x - 8$$

$$\rightarrow y = 2^0 - 8$$

$$\rightarrow 8 = 2^3 = 2^x$$

$$\rightarrow y = 1 - 8$$

$$\rightarrow \therefore x = 3$$

$$\rightarrow y = -7$$

$$\therefore A(3; 0) \quad \checkmark$$

$$\therefore B(0; -7) \quad \checkmark$$

- (2) Write down the equation of the asymptote of graph f .

(1)

$$y = -8 \quad \checkmark$$

- (3) Determine the equation of h if $h(x) = f(2x) + 8$.

(2)

$$h(x) = f(2x) + 8 = 2^{2x} - 8 + 8 \rightarrow h(x) = 2^{2x} = 4^x \quad \checkmark \checkmark$$

- (4) Determine the equation of h^{-1} in the form $y = \dots$

(2)

$$h^{-1}: \quad x = 4^y \rightarrow y = \log_4 x \quad \checkmark \checkmark$$

- (5) Write down the equation of p , if p is the reflection of h^{-1} in the x-axis.

(1)

$$p(x) = -\log_4 x \quad \text{or} \quad p(x) = \log_{\frac{1}{4}} x \quad \checkmark$$

(6) Calculate $\sum_{k=0}^3 g(k) - \sum_{k=4}^5 g(k)$. Show ALL calculations. (4)

$$= [g(0) + g(1) + g(2) + g(3)] - [g(4) + g(5)] \quad \checkmark$$

$$= g(0) + g(1) + g(2) + g(3) - g(4) - g(5) \quad \text{with } x = 3 \text{ the line of symmetry}$$

$$= g(0) + g(3) \quad \checkmark \quad \text{because } g(1) = g(5) \quad \text{and} \quad g(2) = g(4)$$

$$= 4, 5 - 0 \quad \checkmark \quad \rightarrow \quad \text{Given } C(0 ; 4, 5) \quad \text{and calculated in (a) } A(3 ; 0)$$

$$= 4, 5 \quad \checkmark$$